Brazilian Journal of Poultry Science
Revista Brasileira de Ciência Avícola

ISSN 1516-635X 2023 / v.25 / n.4 / 001-008

http://dx.doi.org/10.1590/1806-9061-2023-1784

Original Article

Orange Peel Oil Supplementation in Laying Quail Diet: Effects on Performance and Oxidative Stability of Egg Yolk

■Author(s)

Eratak S[I] https://orcid.org/0000-0003-3788-8704
Guvenc Bayram G[II] https://orcid.org/0000-0002-1413-3651
Semen Z[III] https://orcid.org/0000-0002-7722-5772
Cabuk M[IV] https://orcid.org/0000-0002-5962-1958

[I] Department of Poultry Science, Akhisar Vocational School, Manisa Celal Bayar University, Manisa 45200, Turkey.
[II] Department of Physiology, Faculty of Veterinary Medicine, Dokuz Eylul University, 35890, Izmir, Turkey.
[III] Department of Biochemistry, Faculty of Veterinary Medicine, Dokuz Eylul University, 35890, Izmir, Turkey.
[IV] Department of Animal Nutrition and Nutritional Diseases, Faculty of Veterinary Medicine, Dokuz Eylul University, 35890, Izmir, Turkey.

■Mail Address

Corresponding author e-mail address
Metin Cabuk
Department of Animal Nutrition and Nutritional Diseases, Faculty of Veterinary Medicine, Dokuz Eylul University, 35890, Izmir, Turkey.
Phone: +90 232 572 35 30 / 14504
Email: metin.cabuk@deu.edu.tr

■Keywords

Egg, Japanese quail, Orange peel essential oil, Performance, Malondialdehyde.

CC BY

Submitted: 10/March/2023
Approved: 31/July/2023

ABSTRACT

The present study aimed to investigate the potential utilization of orange peel essential oil (OPEO), as a natural feed additive in poultry nutrition. The impact of incorporating OPEO into the diets of laying quails was examined with respect to various performance criteria, including feed consumption and efficiency, as well as egg production, weight, and shelf life. A total of 120 female egg quails, aged 16 weeks, were allocated into three main groups, and fed with control rations supplemented with OPEO at levels of 0 mg/kg (control group), 200 mg/kg, and 300 mg/kg. Throughout the study, egg production was monitored on a daily basis, feed consumption was recorded every two weeks, and egg weights were measured twice a week on consecutive days. Moreover, the levels of malondialdehyde, a lipid peroxidation marker, in the egg yolks were quantified. The results indicated that the inclusion of OPEO in quail rations did not exert a significant influence on feed consumption and efficiency. However, OPEO supplementation had a positive impact on egg production and weight, leading to substantial increases in both crucial performance parameters. Furthermore, the study elucidated a statistically significant impact of OPEO on the levels of malondialdehyde in the egg yolks. In summary, the incorporation of OPEO into quail diets showcased remarkable efficacy in significantly augmenting egg yield and weight, while leaving feed consumption unaffected. These results highlight the potential of utilizing OPEO as a natural feed additive to improve the performance and egg quality of laying quails in commercial production systems.

INTRODUCTION

The global demand for poultry products has been steadily increasing in response to the growing population and changing dietary preferences (FAO, 2013). To meet this rising demand, the development of innovative strategies to enhance animal productivity in the poultry industry has become imperative. Japanese quail (*Coturnix coturnix japonica*) has gained recognition as a model species in the meat and egg production industry due to its distinct characteristics, including small body size, fast growth rate, and short generation interval (Jeke *et al.*, 2018; Elsaidy *et al.*, 2021). Furthermore, Japanese quail farming offers significant advantages in developing countries as it provides a cost-effective source of animal protein, characterized by its low price, high-quality protein content, and economic value. However, intensive quail rearing practices can pose challenges, including reduced animal immunity, which in turn leads to decreased meat and egg production. Traditionally, the poultry industry has relied on the use of antibiotic growth promoters to mitigate such challenges. However, since the European Union banned the use of antibiotics in poultry feeding in

2006 due to concerns regarding antibiotic resistance and public health, there has been a growing need to explore alternative feed additives that can improve productivity and maintain animal health (EC, 2006). Consequently, researchers and industry professionals have shifted their focus towards finding effective and sustainable alternatives to antibiotic growth promoters (Gong *et al.*, 2014; Zeng *et al.*, 2015; Jamil *et al.*, 2022).

Fruit and vegetable processing methods generates substantial amounts of waste and by-products, resulting in environmental pollution and economic losses (Domínguez *et al.*, 2020). The principles of the circular economy advocate for the utilization of renewable biological resources to produce value-added products, including food and animal feed (Georganas *et al.*, 2020). Notably, vegetable and fruit processing residues contain a wide array of bioactive compounds that offer potential health benefits (Costa *et al.*, 2017). Consequently, their incorporation into animal diets not only addresses waste management challenges but also provides an avenue to develop value-added products (Truong *et al.*, 2019). Among the various residues, citrus peels and components derived from the fruit juice industry have emerged as viable candidates for evaluation as functional compounds. Citrus essential oils derived from citrus peels exhibit a broad spectrum of biological activities, including antimicrobial, antifungal, antioxidant, anti-inflammatory, and anxiolytic properties, making them promising candidates as food and feed additives (Singh *et al.*, 2021).

Previous studies have demonstrated the beneficial effects of essential oil mixtures containing citrus peel oils in broiler chickens, showing increased body weight and feed efficiency when compared to antibiotic-treated groups (Alçiçek *et al.*, 2003). Similar positive outcomes have been observed in layer quails, with improved egg yield and feed efficiency upon the addition of essential oil mixtures containing citrus peel oils (Cabuk *et al.*, 2014).

Among the citrus essential oils, orange peel essential oil (OPEO) is particularly noteworthy. OPEO is concentrated in the colored (flavedo) part of the peel and pigment glands, constituting approximately 0.2-0.5% of the whole fruit. The primary component of OPEO, "limonene" (D-Limonene, 1-methyl-4-(1-methylethenyl) cyclohexane), represents approximately 95% of the oil and is considered safe for use in various industries, including food, cosmetics, perfumery, gum, beverages, and soap, and Generally Recognized as Safe (GRAS) additive (Hosni *et al.*, 2010; Farhat *et al.*, 2011).

While OPEO has recently been utilized as an additive in poultry rations, research investigating its effects is still limited. Previous studies have shown that the addition of OPEO to broiler diets at different concentrations resulted in increased body weight and feed conversion efficiency, with a positive correlation between OPEO concentration and performance (Aydın & Alçiçek, 2018). Furthermore, supplementing OPEO in the diet of heat-stressed chickens has been found to increase egg weight and shell thickness (Özek, 2012), while also improving egg quality in layer quails (Erisir *et al.*, 2015). Additionally, OPEO has been shown to positively impact serum parameters in both chickens (Alzawqari *et al.*, 2016) and layer quails (Sevim *et al.*, 2020).

In light of the extensive knowledge presented, the primary focus of this study was to investigate the impact of incorporating orange peel essential oil (OPEO), derived from the by-product of the fruit juice industry, into the feed of quails – an important poultry species raised for its meat and egg production. Specifically, this research aimed to comprehensively examine the effects of OPEO supplementation on key performance indicators of laying quails, including egg production, egg weight, feed consumption, feed efficiency, and egg shelf life. By conducting a systematic evaluation of the effects of OPEO on these parameters, this study contributes to a deeper understanding of its potential as a natural feed additive in enhancing the productivity and quality of poultry products.

MATERIALS AND METHODS

Animals

The animal model employed in this study consisted of 120 female laying quails (*Coturnix coturnix japonica*) at 16 weeks of age. The animals were housed under conditions of controlled lighting, following a 16-hour light and 8-hour dark cycle. Throughout the investigation, the quails were provided *ad-libitum* access to both feed and water. The ethical implications of the study were carefully addressed, and all experimental procedures were conducted in accordance with the guidelines set forth by the Manisa Celal Bayar University Animal Experiments Local Ethics Committee, adhering to the National Institutes of Health Guidelines for the Care and Use of Laboratory Animals (Decision No: 77.637.435).

Experimental procedure

A total of 120 female laying quails were divided into four subgroups, with each group consisting of 40 quails. Within each group, subgroups of ten quails were created. The quails were housed in multi-story layer quail cages, ensuring equal representation of each subgroup on each floor of the cages. This setup provided standardized housing conditions throughout the study.

The compound feed used in the laying quail trials was formulated with specific raw materials and nutrient compositions, as outlined in Table 1. The feed composition aimed to provide optimal nutrition for the quails during the study.

Table 1 – Ingredent and nutrient composition of the basal experimental layer quail diet (as fed).

Ingredent	
Corn	57.73
Soybean meal (0.48 CP)	19.18
Sunflower meal (0.37 CP)	8.33
Fish meal (0.70 CP)	2.00
Vegetable Oils	3.90
Ground limestone	6.96
Monocalcium phosphate	1.14
Salt	0.20
Vitamin-Mineral premix*	0.25
DL-methionine	0.32
Total	100.00
Nutrient Composition, %	
Dry matter	89.56
Crude Protein (CP)	18.00
Crude fiber	3.25
Crude ash	10.74
Ether extract	6.14
Methionine + cystine	0.82
Lysine	0.88
Calcium	3.00
Phosphorus, available	0.60
Metabolisable Energy, kcal/kg	2900

*Vitamin-mineral premix in 3.0 kg: Vitamin A 12000000 IU; Vitamin D3 3000000 IU; Vitamin E 30000 mg; Vitamin K3 5000 mg; Vitamin B1, 3000 mg; Vitamin B2 12000 mg; Vitamin B6 4000 mg; Vitamin B12 30 mg; Nicotine amide 55000 mg; Calcium--D-Pantothenate 15000 mg; Folic acid 2000 mg; D- Biotin 250 mg; Choline Chloride 600000 mg; Antioxidant 10000 mg; manganese 80000 mg; Iron 40000 mg; Zinc 60000 mg; Copper 5000 mg; Iodine 400 mg; Cobalt 100mg; Selenium 150mg.

The quails were divided into three main groups and fed three different diets throughout the experimental period. The groups included: (1) Control diet without orange peel essential oil (OPEO) (basal diet), (2) Basal diet with 200 mg/kg OPEO, and (3) Basal diet with 300 mg/kg OPEO. OPEO was mixed with a carrier and added at a rate of 1 kg per ton of basal diet. Care was taken to ensure thorough mixing to achieve homogenous distribution of OPEO in the ration.

Throughout the 8-week experimental period, the quails' egg production was recorded daily. Feed consumption and feed efficiency were assessed by weighing the feed every two weeks. Additionally, the weight of the eggs was measured twice a week on consecutive days, with a precision of 0.1 grams. Egg weights for the experimental groups were recorded starting from the second week of the experiment. Eggs were collected on two consecutive days each week for this purpose.

At the end of the 8th week of the experiment, 16 randomly selected eggs per diet treatment (4 from each subgroup) were stored at a temperature of +4°C for durations of 1, 15, and 30 days. To evaluate indicators of fatty acid oxidation and staling criteria, the levels of malondialdehyde (MDA) were measured. A total of 48 egg yolks, representative of each diet group, were used for MDA analysis. The concentration of MDA, an index of lipid peroxidation, was determined using the thiobarbituric acid reaction method, as described by Ohkawa *et al.* (1979).

Determination of Thiobarbituric Acid Reactive Substance (TBARS) Value

To determine the Thiobarbituric Acid Reactive Substance (TBARS) value, a TCA solution (6 ml) consisting of 7.5% TCA, 0.1% EDTA, and 0.1% Propyl gallate (dissolved in 3 ml ethanol) was added to each collected egg yolk. The samples were homogenized using a high-speed polytron for 15 seconds and subsequently filtered through Whatman No:1 filter paper. From the filtrate, 1 ml was transferred to a test tube, followed by the addition of 1 ml of 0.02 M TBA solution. The contents of the test tubes, which were glass test tubes with screw caps, were thoroughly mixed. The test tubes were then placed in a water bath at 100°C for 40 minutes, ensuring proper heating. Afterward, the tubes were promptly cooled in cold water for 5 minutes. Subsequently, the tubes were subjected to centrifugation at 2000 rpm for 5 minutes. The absorbance of the resulting supernatant was measured at a wavelength of 530 nm using a spectrophotometer. The TBARS value, expressed as mg malondialdehyde (MDA) per kg, was determined based on the absorbance readings obtained from the spectrophotometer. This measurement served as an indicator of lipid peroxidation in the egg yolks.

Statistical Analyses

The data obtained from the experiment were subjected to statistical analysis using the General Linear Model procedure implemented in the SAS software (SAS

Institute, 1991). To determine significant differences between the experimental groups, Duncan's multiple range tests were employed. A significance level of $p<0.05$ was considered as statistically significant.

RESULTS AND DISCUSSION

Feed Consumption, Feed Efficiency, and Body Weight

The influence of OPEO on feed consumption, feed efficiency, and final body weight of laying quails is summarized in Table 2. The daily feed consumption of quails in the control group was 30.75 g/quail, while it was 31.16 g/quail in the OPEO (200 mg/kg) group and 30.48 g/quail in the OPEO (300 mg/kg) group. These findings are consistent with previous studies in layer quails (Sevim *et al.*, 2020) and laying hens (Chowdhury *et al.*, 2008) that also reported no significant effect on feed consumption upon OPEO supplementation. However, Erisir *et al.* (2015) observed reduced feed consumption in layer quails supplemented with OPEO. The variation in results could be attributed to the OPEO content and dosage used in the studies, as well as the potential bitter taste of flavonoids in orange peel, which may affect the birds' appetite (Mason *et al.*, 1989; Burdock, 1997).

Table 2 – Effects of OPEO on feed consumption, feed efficiency and body weight in quails.

Treatments	Parameters		
	Feed Intake (g/quail)	Feed Efficiency (g feed/g egg)	Final Body Weight (g)
Control (OPEO-free)	30.75	2.51	279.80
OPEO 200 mg/kg	31.16	2.42	283.23
OPEO 300 mg/kg	30.48	2.45	283.35
SEM Pooled	0.37	0.03	0.363
Probability	0.4357	0.0886	0.4648

OPEO, Orange peel essential oil.

The feed efficiency value, indicating the amount of feed consumed per unit of egg production, improved with the addition of OPEO to the ration, although the difference was not statistically significant. The feed efficiency values were 2.51 g feed/g egg in the control group, 2.42 g feed/g egg in the OPEO (200 mg/kg) group, and 2.45 g feed/g egg in the OPEO (300 mg/kg) group. These results are consistent with other studies that reported no significant change in feed efficiency with OPEO supplementation (Bozkurt *et al.*, 2012; Karabayir *et al.*, 2018; Sevim *et al.*, 2020). Conversely, studies conducted by Bozkurt *et al.* and Çabuk *et al.* (2006) demonstrated that adding an essential oil mixture to laying chicken feeds increased feed efficiency. These findings suggest that the impact of OPEO on feed efficiency may vary across species and depend on the specific formulation and dosage used.

There were no statistically significant effects of OPEO-containing feeds on the final body weight of the quails. The final body weights of quails were 279.80 g in the control group, 283.23 g in the OPEO (200 mg/kg) group, and 283.35 g in the OPEO (300 mg/kg) group. These findings align with previous studies that reported no significant changes in body weight with the addition of orange oil (Karabayir *et al.*, 2018; Sevim *et al.*, 2021) or orange-peel powder (Yildiz *et al.*, 2023) to the ration. The lack of significant effects on body weight suggests that OPEO supplementation within the studied doses does not exert substantial influence on the growth performance of laying quails.

In conclusion, feeding layer quails with diets containing different doses of OPEO did not significantly impact performance parameters. To improve performance quality, further research can explore broader dosage ranges of OPEO or combinations with different essential oil blends to enhance their potential healing effects.

Egg Production and Egg Weight

Significant effects were observed in egg production and egg weight due to the addition of OPEO to the layer quail diet. Supplementation of OPEO in the layer quail diet significantly increased egg production ($p<0.05$) and egg weight ($p<0.01$), as presented in Table 3. Egg production rates were 90.40% in the control group, 92.41% in the OPEO (200 mg/kg) group, and 90.88% in the OPEO (300 mg/kg) group. These findings are consistent with studies reporting positive effects of different doses of OPEO on egg production in poultry (Sevim *et al.*, 2020). Similarly, the addition of an essential oil mixture to the diets of laying chickens has been shown to increase egg production (Çabuk *et al.*, 2006; Bozkurt *et al.*, 2012). However, Özek *et al.* (2011) reported no increase in egg production with the addition of essential oil to hen diets.

Table 3 – Effects of OPEO on egg production and egg weight in quails.

Treatments	Parameters	
	Egg Production (%)	Egg Weight (g)
Control (OPEO-free)	90.40^{b}	12.25^{c}
OPEO 200 mg/kg	92.41^{a}	12.89^{a}
OPEO 300 mg/kg	90.88^{b}	12.45^{b}
SEM Pooled	0.55	0.04
Probability	0.0250	0.0001

^{a,b,c} Means with in column various superscript differ at $p<0.05$ for Egg Production and $p<0.0001$ for Egg Weight.

Egg weight was 12.25 g in the control group, 12.89 g in the OPEO (200 mg/kg) group, and 12.45 g in the OPEO (300 mg/kg) group. Although the 300 mg/kg OPEO dose resulted in slightly lower egg weight compared to the 200 mg/kg dose, it still yielded higher egg weight compared to the control group. Consistent with our findings, Bozkurt *et al.* (2012) reported increased egg weight with the addition of an essential oil mixture to laying hen diets. However, Karabayir *et al.* (2018) found a dose-dependent decrease in egg weight with OPEO supplementation, while Sevim *et al.* (2020) reported no effect on egg weight. These variations in results may be attributed to the specific doses and formulations used in the studies.

Malondialdehyde (MDA) Levels

The effect of OPEO supplementation on the oxidative stability of stored eggs was assessed by measuring malondialdehyde (MDA) levels, a marker of lipid peroxidation. The addition of OPEO in poultry rations resulted in improved egg quality, as evidenced by decreased MDA. Figure 1 illustrates the MDA levels in eggs stored for 1, 15, and 30 days. The MDA accumulation in the control and experimental groups did not differ significantly on the 1st day of storage. However, on the 15th and 30th days, the MDA accumulation in the egg yolk significantly decreased (p<0.0001). For instance, on the 15th day, the MDA accumulation in the control group was 0.179 mg/kg, while it was 0.153 mg/kg in the 200 mg/kg OPEO group and 0.147 mg/kg in the 300 mg/kg OPEO group. Similarly, on the 30th day, the MDA accumulation was 0.274 mg/kg in the control group, 0.207 mg/kg in the OPEO (200 mg/kg) group, and 0.180 mg/kg in the OPEO (300 mg/kg) group.




Figure 1 – The effect of OPEO on the accumulation of MDA (mg/kg yolk) in quail egg yolk during storage (1st, 15th, and 30th days). [a,b,c] means with in column varies superscript differ at p<0.0001.

These results can be attributed to the antioxidant properties of OPEO, which contains phenolic compounds such as Terpinolene and γ-Terpinene known for their inhibitory effects on lipid peroxidation (Ruberto *et al.*, 2000). These compounds reduce the chain length of lipid peroxyl radicals through their active methyl groups, exhibiting antioxidant behavior (Foti & Ingold, 2003; Sharififar *et al.*, 2010). The reduction in MDA levels observed in our study suggests that OPEO, through dietary supplementation, may have prevented the chain reaction involved in the oxidation of dietary lipids, thus enhancing oxidative stability in eggs. These results align with Selcuk & Sengul (2021), who reported decreased MDA values in quails' diets supplemented with lower doses of OPEO. Furthermore, the antioxidant effects of OPEO in poultry diets have been shown to improve both physiological mechanisms (Akbarian *et al.*, 2015) and the quality of animal-derived products (Bolukbasi-Aktas, 2017), supporting our study's results. The preservation of egg quality through the reduction of lipid peroxidation products further supports the potential application of OPEO as a natural antioxidant in the poultry industry.

CONCLUSIONS

In conclusion, our study highlights the significant contributions of orange peel essential oil (OPEO) as a natural additive in layer quail feeds. The addition of OPEO resulted in notable improvements in key yield characteristics such as egg production, egg weight, and egg shelf life. These findings demonstrate the potential of OPEO as a viable substitute for antibiotics in poultry nutrition, providing a natural and herbal growth promoter without the potential harmful residues associated with antibiotic use. Furthermore, the utilization of OPEO, a by-product of the juice industry, aligns with the principles of the circular economy, offering additional environmental and economic benefits. However, further research is necessary to investigate the effects of incorporating OPEO into animal feeds in different combinations and mixtures with other beneficial plant-based essential oils. These studies will enhance our understanding of the synergistic effects and optimal dosages to maximize the benefits of OPEO in poultry production.

By utilizing OPEO as a natural feed additive, the poultry industry can enhance productivity and product quality while reducing reliance on synthetic growth promoters. This research contributes to the development of sustainable and environmentally

friendly strategies in poultry nutrition and supports the shift towards more natural and healthy production practices.

In conclusion, the findings of this study highlight the potential of OPEO as a promising feed additive in layer quail nutrition, promoting the sustainable production of high-quality poultry products.

ACKNOWLEDGMENTS

This study was supported by the Commission of Scientific Research Projects of Manisa Celal Bayar University (Project No: 2019-018).

CONFLICT OF INTEREST

The authors declare that there is no conflict of interests regarding the publication of this article.

REFERENCES

Akbarian A, Golian A, Kermanshahi H, *et al.* Antioxidant enzyme activities, plasma hormone levels and serum metabolites of finishing broiler chickens reared under high ambient temperature and fed lemon and orange peel extracts and Curcuma xanthorrhiza essential oil. Journal of Animal Physiology and Animal Nutrition 2015;99:150-62. https://doi.org/10.1111/jpn.12188

Alçiçek A, Bozkurt M, Çabuk M. The Effect of an essential oil combination derived from selected herbs growing wild in Turkey on broiler performance. South African Journal of Animal Science 2003;33:89-94. http://dx.doi.org/10.4314/sajas.v33i2.3761

Alzawqari MH, Al-Baddany AA, Al-Baadani HH, *et al.* Effect of feeding dried sweet orange (Citrus sinensis) peel and lemon grass (Cymbopogon citratus) leaves on growth performance, carcass traits, serum metabolites and antioxidant status in broiler during the finisher phase. Environmental Science and Pollution Research 2016;23:17077-82. https://doi.org/10.1007/s11356-016-6879-7

Aydın A, Alçiçek A. Effects of the supplementation of essential oil isolated from orange peel (Citrus sinensis L.) to broiler diets on the performance. Türk Tarım ve Doğa Bilimleri Dergisi 2018;5:127-35. https://doi.org/10.30910/turkjans.421348

Bozkurt M, Kucukyılmaz K, Pamukcu M, *et al.* Long-term effects of dietary supplementation with an essential oil mixture on the growth and laying performance of two-layer strains. Italian Journal of Animal Science 2012;11:23-8. https://doi.org/10.4081/ijas.2012.e5

Burdock GA. Encyclopedia of food and color additives. Boca Raton: CRC Press; 1997.

Çabuk M, Bozkurt M, Alçiçek A, *et al.* Effect of dietary essential oil mixture on performance of laying hens in summer season. South African Journal of Animal Science 2006;36:215-21.

Cabuk M, Eratak S, Alçiçek A, *et al.* Effects of herbal essential oil mixture as a dietary supplement on egg production in quail. The Scientific World Journal 2014;573470-4. https://doi.org/10.1155/2014/573470

Chowdhury SD, Hassin BM, Das SC, *et al.* Evaluation of marigold flower and orange skin as sources of xanthophyll pigment for the improvement of egg yolk color. The Journal of Poultry Science 2008;45:265-72. https://doi.org/10.2141/jpsa.45.265

EC - Commission Regulation. Regulation (EC) No. 1831/2003 of the European Parliament and of the Council on additives for use in animal nutrition. Official Journal L 2003;268:29-43.

Costa APD, Hermes VS, Rios AO, *et al.* Minimally processed beetroot waste as an alternative source to obtain functional ingredients. Journal of Food Science and Technology 2017;54:2050-8. https://doi.org/10.1007%2Fs13197-017-2642-4

Domínguez R, Munekata PES, Pateiro M, *et al.* Red beetroot. A potential source of natural additives for the meat industry. Applied Sciences 2020;10:8340. https://doi.org/10.3390/app10238340

Elsaidy N, Kirella A, El-Kassas S, *et al.* Reducing the abundance of harmful bacteria of rooftop tank–stored drinking water using silver nanoparticles and acetic acid and its impact on Japanese quail growth performances. Biological Trace Element Research 2021;199:3062-72. https://doi.org/10.1007/s12011-020-02422-2

Bolukbasi-Aktas SC. Narenciye kabuk yağlarının etlik piliçlerde doku yağ asidi kompozisyonu ve raf ömrü üzerine etkileri. Atatürk Üniversitesi Veteriner Bilimleri Dergisi 2017;12:157-66. https://doi.org/10.17094/ataunivbd.347682

Erisir Z, Şimşek ÜG, Çiftçi M, *et al.* The effect of orange peel oil and sex ratio on egg production and egg characteristics in layer quail (Coturnix coturnix japonica). Fırat University Journal of Health Sciences 2015;29:23-30.

FAO - Food and Agriculture Organization. Poultry development review. Rome; 2013.

Farhat A, Fabiano-Tixier AS, Maataoui ME. Microwave steam diffusion for extraction of essential oil from orange peel: Kinetic data, extract's global yield and mechanism. Food Chemistry 2011;125:255-61. https://doi.org/10.1016/j.foodchem.2010.07.110

Foti MC, Ingold KU. Mechanism of inhibition of lipid peroxidation by gamma-terpinene, an unusual and potentially useful hydrocarbon antioxidant. Journal of Agricultural Food 2003;51:2758-65. https://doi.org/10.1021/jf020993f

Georganas A, Giamouri E, Pappas AC, *et al.* Bioactive compounds in food waste: A review on the transformation of food waste to animal feed. Foods 2020;9:1–18. https://doi.org/10.3390/foods9030291

Gong J, Yin F, Hou Y, *et al.* Review: Chinese herbs as alternatives to antibiotics in feed for swine and poultry production: Potential and challenges in application. Canadian Journal of Animal Science 2014;94:223-41. https://doi.org/10.4141/cjas2013-144

Hosni K, Zahed N, Chrif R, *et al.* Composition of peel essential oils from four selected Tunisian citrus species: Evidence for the genotypic influence. Food Chemistry 2010;123:1098-104. https://doi.org/10.1016/j.foodchem.2010.05.068

Jamil M, Aleem MT, Shaukat A, *et al.* Medicinal plants as an alternative to control poultry parasitic diseases. Life 2022;12:449. https://doi.org/10.3390/life12030449

Jeke A, Phiri C, Chitindingu K, *et al.* Ethnomedicinal use and pharmacological potential of Japanese quail (Coturnix coturnix japonica) birds` meat and eggs, and its potential implications on wild quail conservation in Zimbabwe: a review. Cogent Food & Agriculture 2018;4:1507305. https://doi.org/10.1080/23311932.2018.1507305

Karabayir A, Öğütücü M, Acar Ü, *et al.* Effects of orange peel oil on quail (coturnix coturnix japonica) growth-performance, egg quality and blood parameters. New Knowledge Journal of Science 2018;7(2):127-136.

Mason JR, Adams MA, Clark L. Anthranilate repellency to starlings: chemical correlates and sensory perception. The Journal of Wildlife Management 1989;53:55–64. https://doi.org/10.2307/3801306

Ohkawa H, Ohishi N, Yagi K. Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. Analytical Biochemistry 1979;95:351-358. https://doi.org/10.1016/0003-2697(79)90738-3

Özek K, Wellmann KT, Ertekin B, *et al.* Effects of dietary herbal essential oil mixture and organic acid preparation on laying traits, gastrointestinal tract characteristics, blood parameters and immune response of laying hens in a hot summer season. Journal of Animal and Feed Sciences 2011;20:575-86. https://doi.org/10.22358/jafs/66216/2011

Özek K. Effects of dietary herbal essential oil mixture and/or mannan-oligosaccharide supplementation on laying performance, some serum biochemical markers and humoral immunity in laying hens exposed to heat. Revue de Medecine Veterinaire 2012;163:153-9.

Ruberto G, Baratta MT, Deans SG, *et al.* Antioxidant and antimicrobial activity of Foeniculum vulgare and Crithmum maritimum essential oils. Planta Medica 2000;66:687-93. https://doi.org/10.1055/s-2000-9773

SAS Institute. SAS user's guide: statistics. Version 6.03 Cary: SAS Institute; 1991.

Selcuk S, Sengul T. Portakal kabuğu yağı ve nar çekirdeği yağı ile zenginleştirilen diyetlerin bıldırcınların verim performansı, yumurta kalitesi ve bazı kan parametreleri üzerine etkileri. Türk Tarım ve Doğa Bilimleri Dergisi 2021;8:702-19. https://doi.org/10.30910/turkjans.899877

Sevim B, Gökmen SA, Cufadar Y, *et al.* The effect of addition of orange and lemon peel essential oils and mixtures to breeder quail rations on performance, egg quality and hatch parameters. Turkish Journal of Agriculture - Food Science and Technology 2021;9:1988-92. https://doi.org/10.24925/turjaf.v9i11.1988-1992.4578

Sevim B, Olgun O, Şentürk ET, *et al.* The effect of orange peel oil addition to laying quail diets on performance, eggshell quality and some serum parameters. Turkish Journal of Agriculture -Food Science and Technology 2020;8:1773-7. https://doi.org/10.24925/turjaf.v8i8.1773-1777.3508

Sharififar F, Moshafi MH, Dehghan-Nudehe G, *et al.* Bioassay screening of the essential oil and various extracts from 4 spices medicinal plants. Pakistan Journal of Pharmaceutical Sciences 2009;22:317-22.

Singh B, Singh JP, Kaur A, *et al.* Insights into the chemical composition and bioactivities of citrus peel essential oils. Food Research International 2021;143:110231. https://doi.org/10.1016/j.foodres.2021.110231

Truong L, Morash D, Liu Y, *et al.* Food waste in animal feed with a focus on use for broilers. International Journal of Recycling Organic Waste in Agriculture 2019;8:417-29. https://doi.org/10.1007/s40093-019-0276-4

Yildiz A, Kılınc G, Olgun O, *et al.* Effect of supplementation of orange peel powder to the diet on performance, egg quality and egg yolk antioxidant activity in laying quails. Turkish Journal of Agriculture - Food Science and Technology 2023;11:151-55. http://doi.org/10.24925/turjaf.v11i1.151-155.5808

Zeng Z, Zhang S, Wang H, *et al.* Essential oil and aromatic plants as feed additives in non-ruminant nutrition: a review. Journal of Animal Science and Biotechnology 2015;6:7. https://doi.org/10.1186/s40104-015-0004-5